

Skin Rejuvenation Using Intense Pulsed Light ***A Randomized Controlled Split-Face Trial With Blinded Response Evaluation***

Lene Hedelund, MD; Eva Due, MD; Peter Bjerring, MD, DMSc; Hans Christian Wulf, MD, DMSc; Merete Haedersdal, MD, PhD, DMSc

Objective: To evaluate efficacy and adverse effects of intense pulsed light rejuvenation in a homogeneous group of patients.

Design: Randomized controlled split-face trial.

Setting: University dermatology department.

Patients: Thirty-two female volunteers with Fitzpatrick skin type I through III and class I or II rhytids.

Interventions: Subjects were randomized to 3 intense pulsed light treatments at 1-month intervals or to no treatment of right or left sides of the face.

Main Outcome Measures: Primary end points were skin texture and rhytids. Secondary end points were telangiectasia, irregular pigmentation, and adverse effects. Efficacy was evaluated by patient self-assessments and by blinded clinical and photographic valuations up to 9 months after final treatment. Adverse effects were assessed clinically and by noninvasive skin reflectance measurements.

Results: Skin texture was significantly improved at all clinical assessments except at the 6-month examination ($P_{.006}$). The improvements peaked at 1 month after treatment, at which time 23 (82%) of 28 patients had better appearances of treated vs untreated sides. Most patients obtained mild or moderate improvements, and 16 patients (58%) self-reported mild or moderate efficacy on skin texture. Rhytids were not significantly different on treated vs untreated sides, and 19 patients (68%) reported uncertain or no efficacy on rhytids. Significant improvements of telangiectasia ($P_{.001}$) and irregular pigmentation ($P_{.03}$) were found at all assessments. Three patients withdrew from the study because of pain related to treatment.

Conclusions: Three intense pulsed light treatments improved skin texture, telangiectasia, and irregular pigmentation but had no efficacy on rhytids. Adverse events were minimal, but included scar in 1 patient.